




Advancing the RIDBC Renwick Centre experience.

Unified English Braille Training Manual

Extension Mathematics

Josie Howse



 **UEBOnline**

Unified English Braille Training Manual: Extension Mathematics
Revision 4

© 2020 North Rocks Press

NextSense

361-365 North Rocks Road

North Rocks NSW 2151

Australia

<https://www.nextsense.org.au/>

This work is licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International Licence.

- Attribution: You must give appropriate credit to the author
- Non-Commercial: You must not use the material for commercial purposes
- No derivatives: If you remix, transform or build upon the material, you may not distribute the modified material.

To view a copy of this licence, visit <https://creativecommons.org/licenses/by-nc-nd/4.0/> or send a letter to Creative Commons, 171 Second Street, Suite 300, San Francisco, California 94105 USA



ISBN 978-0-949050-07-6



A catalogue record for this book is available from the National Library of Australia

NextSense Institute is Australia's leading centre for research and professional studies in the field of education for children with sensory disabilities; offering webinars, short courses, and degree programs for parents, carers, educators, and health professionals. NextSense Institute is committed to providing high-quality teaching and learning opportunities. Our programs are conducted by leading national and international experts for education and health professionals who support people who are deaf, hard of hearing, blind or have low vision.

Table of Contents

Foreword.....	6
Contributors	8
Lesson 1 - Revision	9
Grade 1 Mode	9
Grade 1 Symbol Indicator	10
Termination of Grade 1 Mode.....	11
Numeric Mode	11
Termination of Numeric mode	12
Standing Alone Rule.....	12
Grade 1 Word Indicator	13
Grade 1 Passage Indicator	14
Grade 1 Passage Terminator	14
Grade 1 Passage Indicator and Terminator on lines of their own	14
Shape Indicators	15
Transcriber defined shapes	16
Use of the shape termination indicator	16
Omissions in technical expressions.....	17
Fractions	17
Simple numeric fraction	17
General Fractions	18
Miscellaneous Symbols.....	21
Superscripts and Subscripts	25
Definition of an Item.....	25
Exercise 1	29
Extra Exercise 1	29
Lesson 2 - Functions	31
Trigonometric functions	31
Exercise 2	35
Review Exercise 2.....	35
Lesson 3 - Modifiers, Bars and Dots.....	37

Modifiers.....	37
Modifiers directly above or below	37
Definition of an Item.....	37
Braille Grouping Symbols	38
Bars	38
Two indicator bars applied to the same item	39
Dots.....	39
Recurring decimal:.....	40
Derivatives of x	40
Derivatives of x	40
Exercise 3	43
Review Exercise 3.....	43
Lesson 4 - Modifiers, Arrows and Limits	45
Modifiers.....	45
Modifiers directly above or below	45
Definition of an Item.....	45
Braille Grouping Symbols	46
Arrows	47
Simple Arrows	47
Arrow symbol directly above and below	48
The Limit Function.....	49
Exercise 4	50
Review Exercise 4.....	51
Lesson 5 - Greek Letters (continued) and Probability	52
Greek Letters	52
Probability	54
Venn Diagrams	55
Complementary Events	56
Independent Events.....	56
Dependent Events	57
Exercise 5	59

Review Exercise 5.....	59
Lesson 6 - Calculus: Differentiation	61
Exercise 6	64
Review Exercise 6.....	64
Lesson 7 - Calculus: Integration	66
Exercise 7	68
Review Exercise 7.....	68
Lesson 8 - Vertical Bars.....	70
Exercise 8	72
Review Exercise 8.....	72
Lesson 9 - Review Test	74
Review Test	76
Lesson 10 - Extension Test	78
Extension Test	80
References	82

Foreword

We live in an era where education leaders and decision makers are responding to national and international disability anti-discrimination legislation, policies and standards that prioritise equitable inclusive education for all, leaving no-one behind. However, for learners with vision impairment, the low incidence nature of their disabilities can present challenges for education systems and schools. The World Blind Union in 2019 highlighted three global challenges impeding education enrolment and achievement for children and young people with vision impairment, in particular in low and middle-income countries. These challenges are a chronic shortage of teachers who are qualified to teach braille, an absence of braille instruction in teacher training programs or offered by education systems, and the high cost of braille equipment. With these challenges in mind, NextSense has created online, open access training programs and resources in braille literacy and mathematics for education leaders, teachers, parents and caregivers – basically, for anyone who wants to learn braille.

In my experience as a lecturer in sensory impairment, professional qualifications, quality teaching and positive student outcomes are closely linked. Knowledgeable and inspiring teachers reach out to their students, building and enriching student knowledge and igniting curiosity, self-confidence, and motivation to pursue learning. In the field of mathematics, we need such teachers who can reach out to their students with vision impairment. Teacher training institutions have a responsibility to provide teachers with instruction in braille codes and braille literacy development, the mechanics of producing braille and tactile representations of visual information, and practical experience in reading and transcribing print and braille materials.

In offering this UEB Extension Mathematics training manual and online program, NextSense **aims** to address the following:

- To extend professional and parent/caregiver knowledge of braille mathematics by building upon the lessons and exercises presented in the introductory and advanced braille mathematics training programs
- To introduce mathematical symbols in braille that are commonly found in senior secondary mathematics publications; and in so doing,
- To support and promote equitable access to mathematics for learners who use braille or a combination of braille, print and digital formats.

In Australia, as in other parts of the world, braille experts who are knowledgeable in the field of mathematics are limited in number. NextSense is fortunate to have Josie Howse write the content of this Extension Mathematics training manual, in addition to the Introductory and Advanced Mathematics training manuals. During her employment and professional affiliations over the past four decades, Josie Howse has been at the forefront of braille training, assessment, and production in Australia and overseas. She has

contributed to the development of Unified English Braille (UEB) by the International Council on English Braille (ICEB), and adoption and implementation of Unified English Braille by Australia in 2005.

NextSense acknowledges the contribution of Craig Cashmore of Peppacode in developing and maintaining the UEB Online website and translating the content of the literary and mathematics training manuals into accessible, interactive online training programs. Sincere thanks are also extended to the NextSense UEB Online design team and the foundations and trusts that have contributed to the UEB Online training programs since 2014.

I wish to reflect on a quotation from Hellen Keller, as published by the American Foundation for the Blind. Hellen Keller spoke these words 14 years before the United Nations "Universal Declaration of Human Rights" in 1948. Her words are just as relevant today and serve as a clarion call to us all to continue our efforts to promote and support the right to education for all children, leaving no-one behind.

"Education should train the child to use his brains, to make for himself a place in the world and maintain his rights even when it seems that society would shove him into the scrap-heap."

Hellen Keller, 1934

Frances Gentle, AO, PhD, D. Litt. *Honoris Causa*
Conjoint Lecturer, NextSense Institute,
President, International Council for Education of People with Visual Impairment; and Co-
President, South Pacific Educators in Vision Impairment

Contributors

Author: Josie Howse, PSM

M.Spec. Ed. (Sensory Disability) – University of Newcastle,
BA (Ed.) – Macquarie University

Josie Howse is an Adjunct Research Fellow with the NextSense Institute and former Manager of the Braille and Large Print Services, NSW Department of Education. The NSW Department of Education team provides all texts and examinations in braille, large print and e-text to students with vision impairment in the government sector and is the largest producer of alternate format student textbooks and examinations in Australia.

Josie has been working in the field of vision impairment for more than 40 years. She has held a number of executive positions at national and state levels and has extensive experience in braille code development at an international level. Josie was the editor of the 2006 Unified English Braille Primer: Australian Edition, and co-editor of the 2016 Unified English Braille: Australian Training Manual.

Josie was awarded the Public Service Medal (PSM) in the 2007 Queen's Birthday Honours list, has been listed annually in the Who's Who of Australian Women since 2008, and is the 2012 recipient of a Lifetime Achievement Award from the Round Table on Information Access for People with Print Disabilities, and awarded Honorary Life Membership of the South Pacific Educators in Vision Impaired (SPEVI) in 2020.

UEB Online Developer: Craig Cashmore

B. Eng. (Hons) - UTS

Craig holds an Engineering Degree in Telecommunications and has worked in the software development industry for over 30 years, holding senior software design, software architecture and technical management positions in companies including Jtec, Ericsson and LongReach Networks.

More recently Craig founded Peppacode, a web and app development business focused on 'out-of-the-ordinary' strategic web and software development for small business, start-ups and educational institutions.

Some of Craig's achievements at Peppacode include the successful launch of UEB Online for NextSense and a vehicle tracking and management system for a bus operator. Craig continues to work on new and innovative projects using modern web technologies.

Lesson 1 - Revision

Welcome to the Unified English Braille (UEB) Training Manual: Extension Mathematics. This training program builds on the basic principles that you have already learnt in the Introductory and Advanced level training programs but will apply the principles in an extended mathematics context. You will also be introduced to some new signs which may be required when transcribing higher level mathematics.

All examples and test items presented in this training program have been taken directly from Senior Mathematics textbooks.

Lesson 1 will review some of these critical principles to reinforce what has already been delivered in the Introductory and Advanced mathematics programs and prepare you for implementing best practice in this extension training program.

Some of the topics covered in this revision lesson are as follows:

- Grade 1 Mode
- Shape Indicators
- Fractions
- Miscellaneous Symbols
- Superscripts and Subscripts

Grade 1 Mode

The rules for the use of Grade 1 mode in a literary context will also apply in a mathematical context and are summarised below.

A braille symbol may have both an uncontracted (Grade 1) meaning and a contracted (Grade 2) meaning. A Grade 1 indicator is used to set Grade 1 mode when the Grade 1 meaning of a symbol could be misread as a contraction meaning or a numeric meaning.


The extent of Grade 1 mode is determined by the Grade 1 indicator in use.



Grade 1 Symbol Indicator (dots 5 6)



Grade 1 Word Indicator (dots 5 6, 5 6)



Grade 1 Passage Indicator (dots 5 6, 5 6, 5 6)



Grade 1 Passage Terminator (dots 5 6, 3)



Grade 1 Passage Indicator on a line of its own



Grade 1 Passage Terminator on a line of its own

Note:


- Grade 1 indicators will not be needed for simple arithmetic problems involving numbers, operation signs, numerical fractions, and mixed numbers.

Grade 1 Symbol Indicator

- A Grade 1 symbol indicator (⠠) sets Grade 1 mode for only the next symbol.

Example:


In Jones¹⁵ you will find the reference.



- A Grade 1 symbol indicator is NOT required before the letters a, i and o, because they do not have a contraction meaning when they “stand alone”.

Example:

The letters a, i and o do not need a Grade 1 indicator.



- A Grade 1 symbol indicator precedes a capitalisation indicator.

Example:

The letter B.



- A Grade 1 indicator will be required when a lower-case letter a – j immediately follows a digit so it will not be misread as part of the number.

Examples:

25b ⠠⠠⠠⠠⠠⠠ not 252

Trevor@take5.com



“b”



(b)



(B)



“a’s and b’s”



“A’s and B’s”



“can’t”



Grade 1 Word Indicator

- The Grade 1 word indicator (⠠ ⠠) sets Grade 1 mode for the next symbols-sequence or the remainder of the current symbols-sequence.
- The effect of a Grade 1 word indicator is terminated by a space or a Grade 1 terminator.
- Remember that the numeric indicator also sets Grade 1 mode for the next symbols-sequence.
- If a complex algebraic expression does NOT include a comparison sign (such as an equal's sign) then it is unlikely to include interior spaces in braille. In such cases, a Grade 1 word indicator will be enough to ensure that superscripts, subscripts, fractions, radicals, arrows, and shape indicators are well defined without the need for Grade 1 symbol indicators.


Examples:

- Evaluate $\sqrt{y - x^2}$



2. $x^2 - 4x - 3 = 0$

3. $2x^2 - x = 1$



Shape Indicators

A number of shapes used in mathematics were introduced in the UEB Introductory and Advanced Mathematics training manuals. The most common shapes used in senior mathematics are shown below.

Please note that the shape indicator also has a contracted (Grade 2) meaning and so may require a Grade 1 indicator in front of it. A Grade 1 indicator is used to set Grade 1 mode when the Grade 1 meaning of the symbol could be misread as a contraction meaning and Grade 1 mode has not already been established in the sequence.



shape indicator



shape termination indicator (there is no print representation for this braille symbol)




square



triangle (equilateral)



pentagon



hexagon



octagon

⠠⠠⠠⠠⠠⠠ parallelogram

⠠⠠⠠ circle

⠠ visible space

Transcriber defined shapes

⠠⠠ transcriber-assigned shape indicator

Note: A transcriber-assigned shape indicator should **not** be used if the print symbol has already an assigned braille sign. The indicator should precede a short series of initials or a single Grade 1 word. The definitions of all the transcriber-assigned shape indicators used in the transcription should be available to the reader in either a transcriber's note or on a specific page.

Example:

A smiling face icon could be shown either of the following ways:

⠠⠠⠠⠠⠠⠠ or ⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠

Use of the shape termination indicator

⠠ Shape termination indicator (there is no print representation for this symbol in braille)

- If a shape indicator is followed by a space, then no termination sign is required.
- If the shape indicator is followed by punctuation or is unspaced from a following symbol, then the shape termination indicator must be used.
- All the initial shape indicators initiate shape mode so no further Grade 1 indicators will be needed.

Examples:

Δ ABC

⠠⠠⠠⠠⠠⠠ ⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠

Δ,

⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠

ΔABC

⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠

Omissions in technical expressions

Follow the print when omissions are indicated by shapes (such as squares) or punctuation (such as underscores or question marks).

When print uses a blank within a technical expression, use the visible space indicator (⠠) to represent the blank.

Space the sign of omission the same as the sign being omitted.

Remember a Grade 1 indicator is used to set Grade 1 mode when the Grade 1 meaning of the symbol could be misread as a contraction meaning and Grade 1 mode has not already been established in the sequence.

Examples:


$$\square + 75 = 80$$



$$48 = 12 \times \square + \Delta$$



$$? \div 6 = 84$$



$$50 - \quad = 40$$



Fractions

It is important to understand the definition of a **simple fraction**, which the elements and only those elements that can be considered to be a simple fraction and subsequently use the simple fraction line, (⠠) between the numerator and the denominator.

If a fraction does **not** comply with the **simple fraction** definition (restated below) for whatever reason, then it will be a **general fraction** and will require a quite different approach, using different signs as described in a section that follows.

Simple numeric fraction

A simple numeric fraction is one whose numerator (top of the fraction line) and denominator (bottom of the fraction line) contain only:

$\mathbb{1}$		vertical bar (absolute value)
$\mathbb{2}$	∞	infinity
$\mathbb{3}$!	factorial sign
$\mathbb{4}$	\therefore	therefore sign
$\mathbb{5}$	\sphericalangle	angle sign
$\mathbb{6}$		parallel to
$\mathbb{7}$	\perp	perpendicular to
$\mathbb{8}$	\equiv	is congruent/equivalent to
$\mathbb{9}$		is similar to
$\mathbb{10}$	[open square bracket
$\mathbb{11}$]	close square bracket
$\mathbb{12}$	{	open curly bracket
$\mathbb{13}$	}	close curly bracket
$\mathbb{14}$		dot 5 continuation indicator (positioned up close to the last character when the braille sequence is too long for the line and needs to be broken at a logical place)
$\mathbb{15}$		visible blank space (an omission, which often occurs in fractions)
$\mathbb{16}$	/	cancelling sign
$\mathbb{17}$	✓	tick sign (not to be confused with the root sign)

Notes:

- The spacing of symbols generally follows the print.



Superscripts and Subscripts

⠠	Level change up
⠡	Level change down
⠢	Open braille grouping sign
⠣	Closing braille grouping sign

Definition of an Item

An “item” is defined as any single symbol(s) that follows immediately after the level change indicator. It is therefore important to make clear to the reader exactly what symbol(s) must be included as a consequence of the level change indicator.

There is a defined list below of specific conditions that are considered to be the “next item”.

1. An entire number expressed in braille, i.e. the initiating numeric indicator and all succeeding symbols within the numeric mode (including any interior decimal points, commas, separator spaces, or simple numeric fraction lines).
2. An entire general fraction enclosed in general fraction indicators (⠢⠠) and (⠡⠠). Note that Grade 1 indicators will be required if the sequence is not already in Grade 1 mode.
3. An entire radical expression enclosed in radical indicators (such as a square root).
4. An arrow.
5. An arbitrary shape.
6. Any expression enclosed in matching pairs of round parentheses, square brackets or curly braces.

Notes:

- If none of the above conditions apply then the “item” is only the next symbol and may require braille grouping indicators, opening (⠢) and closing (⠣), to ensure the whole of the superscript or subscript has been captured. Note that Grade 1 indicators will be required for these signs if the sequence is not already in Grade 1 mode.

- Any expression when enclosed in the braille grouping indicators described above will subsequently make it clear to the reader exactly what symbols are included as part of the level change.
- A negative superscript or subscript must be enclosed in braille grouping signs because a minus sign can be an item in its own right.
- The superscript and subscript signs each have a Grade 2 (contracted) meaning so will always require Grade 1 indicators if the sequence is not already shown in Grade 1 mode.
- The numeric indicator also sets Grade 1 mode for the remainder of the symbols-sequence.

Examples:

$$2^4$$




$$x^{4.4}$$



$$x^{\frac{1}{4}}$$



$$x^{\frac{a}{b}}$$



preferred

OR



$$x^{(-5)}$$



$$x^n$$



x^{-n}



x^{-2b}



OR



x_{2b}




OR



x_n




y_{-n}



OR



$$\frac{p_1 + p_2}{p_1 - p_2}$$



Simultaneous Superscripts or Subscripts

- If more than one superscript or subscript applies to the term, work from bottom to top, or left to right.

Examples:

$x_1^2 = y_2^3$



$$x^2_k$$



Left-displaced superscripts or subscripts

- When both a subscript and a superscript are shown together in print to the left of the base symbol, they are brailled by using the corresponding index expression prior to the base symbol, working from bottom to top.
- Note the Grade 1 indicator that is required in front of the subscript at the beginning of the example below but not required for the superscript sign, due to the numeric indicator setting Grade 1 mode for the remainder of that sequence.

Example:

$${}^{238}{}_{92}U$$



General Note: There is often a choice of Grade 1 indicators in mathematical contexts, with use of any of the options for Grade 1 mode (symbol, word or passage) being equally correct.

Decisions about option selection are generally associated with user and transcriber preferences, or local production formatting guidelines, and include consideration for simplicity or functionality.

Therefore, for the purpose of the exercises in this training program please use the following criteria for implementing Grade 1 mode:

- Use the Grade 1 **symbol** indicator when there is only one symbol in the sequence requiring a Grade 1 indicator.
- Use the Grade 1 **word** indicator when there are two or more symbols in the sequence requiring Grade 1 mode except in a context whereby any literary elements will be affected and as a consequence will also be uncontracted.
- Use Grade 1 **passage** indicator (with Grade 1 terminator) when a comparison indicator or a space is used in the sequence and Grade 1 indicators are required on both sides of the equation. Remember, however, the impact of Grade 1 mode on any literary elements.

Exercise 1

1. $\frac{r-3}{2} \leq -6$

2. Simplify $\frac{2}{3} - \frac{x-1}{4}$

3. $\therefore x = \frac{5 + \sqrt{45}}{2}$

4. $q = \sqrt[3]{x^3 + y^3 + z^3}$

5. $\frac{m^2-8}{m+1} \geq 4$

6. $\left(\frac{9}{25}\right)^{k+3} = \sqrt{\frac{3}{5}}$

7. $\frac{y}{w} > \frac{1+\sqrt{5}}{2}$

8. $\frac{42g^3h^4}{7h^2} =$

9. $S = \sqrt{\frac{3V}{h}}$

10. $A = \frac{h}{3}(d_f + 4d_m + d_1)$

Extra Exercise 1

1. $\left(\frac{1}{27}\right)^{3n+1} = \frac{\sqrt{3}}{81}$

2. $A = \frac{2 - \sin\theta}{2 \cos\theta}$

3. $T_n = 5^n - 1$

4. $x = t\sqrt{49 - t^2}$

5. $\log\left(\frac{x^2}{y}\right) = \log 2$

6. $(x - h)^2 = \pm 4a(y - k)$

7. Let $r = \sqrt{\frac{4p}{3}}$

8. $\frac{2b}{5} - \frac{1}{2} \geq 6$

9. $\frac{q-2}{3} < 2 + \frac{3q}{4}$

10. $\left(x + \frac{1}{x}\right)^2$

⠠⠠⠠⠠⠠⠠⠠⠠⠠

- Insert a space if a function name is **preceded** directly by a lower or upper-case Latin letter, with no intervening braille indicators or brackets.

Examples:

xcos 30

⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠

Xcos 30

⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠

- Insert a space if a function name is **followed** directly by a lower-case letter with no intervening braille indicators or brackets.

Example:

log x

⠠⠠⠠⠠⠠⠠⠠⠠

- The space is not needed if the function name is already separated by a bracket or by a braille indicator, such as a capitalisation indicator, a Greek letter indicator or an opening general fraction indicator.
- Remember the opening and closing general fraction indicators have a Grade 2 (contracted) meaning so unless you are already in Grade 1 mode, a Grade 1 symbol indicator will be required before the opening and closing fraction indicator.
- We will investigate Greek letters more thoroughly in Lesson 5 of this Training Manual.

Examples:

log (x+y)

⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠

cos *B*

⠠⠠⠠⠠⠠⠠⠠⠠⠠


Tan θ

⠠⠠⠠⠠⠠⠠⠠⠠⠠

tan $\frac{\theta}{2}$

Example:

$$\text{Lim } f(x) = 1$$



Remember: As referred to in the General Note in Lesson 1, there is often a choice of Grade 1 indicators in mathematical contexts, with any of the options of Grade 1 mode (symbol, word or passage) being equally correct.

Decisions about option selection are generally associated with user and transcriber preferences, or local production formatting guidelines, and include consideration for simplicity or functionality.

Therefore, for the purpose of the exercises in this training program, please use the following criteria for implementing Grade 1 mode:

- Use the Grade 1 **symbol** indicator when there is only one symbol in the sequence requiring a Grade 1 indicator.
- Use the Grade 1 **word** indicator when there are two or more symbols in the sequence requiring Grade 1 mode except in a context whereby any literary elements will be affected and as a consequence will also be uncontracted.
- Use Grade 1 **passage** indicator (with Grade 1 terminator) when a comparison indicator or a space is used in the sequence and Grade 1 indicators are required on both sides of the equation. Remember however, the impact of Grade 1 mode on any literary elements.

Hint:

- The numeric indicator also sets Grade 1 mode for the remainder of the symbols-sequence.

Exercise 2

1. $\text{Log} \frac{y}{4}$
2. $-\frac{1}{4} \sec^4 x$
3. $\text{Log}_2 \frac{\sqrt{2}}{4}$
4. $\theta = \cos^{-1} \left(\frac{5.8}{7.3} \right)$
5. $\ln 9.8 + \log_{10} 17$
6. $\log_a \left(\frac{x}{y} \right)$
7. $\theta = n\pi + \tan^{-1} a$
8. $\log_a x = \frac{\log_b x}{\log_b a}$
9. $\cot A = \frac{\cos A}{\sin A}$
10. $\sec A = \frac{1}{\cos A}$

Review Exercise 2

1. $a^{3y-5} = \frac{1}{a^2}$
2. $\ln(4x + 5)$
3. $y - y_1 = m(x - x_1)$
4. $\sin^{-1} \frac{x}{a} + C$
5. $\log_a x^n = n \log_a x$
6. $K = n \sqrt{\frac{a}{p}}$

7. $\log_a \sqrt{xy}$

8. $f(x) = 3x^4 + 4x^3 - 12x^2$

9. $\log \sqrt{x^2 - 4x + 4} - \log(x - 2)$

10. $\frac{-5 \pm \sqrt{21}}{2}$

Lesson 3 - Modifiers, Bars and Dots

Modifiers

Modifiers directly above or below

If an item is written in print directly above or directly below a term, rather than to the right or left, use the directly above indicator or directly below indicator instead of the superscript or subscript. Further explanation of a Modifier directly above or below is explained in Lesson 4.

As stated below in this lesson, special signs are provided for bars and a dot directly above and directly below.

⠠⠠⠠ Expression directly above

⠨⠠⠠⠠ Expression directly below

Definition of an Item

An “item” is defined as any single symbol(s) that follows immediately after the level change indicator. It is therefore important to make clear to the braille reader exactly what symbol(s) must be included as a consequence of the level change indicator.

Below is a defined list of specific conditions that are considered to be the “next item”:

- An entire number expressed in braille, i.e., the initiating numeric indicator and all succeeding symbols within the numeric mode (including any interior decimal points, commas, separator spaces, or simple numeric fraction lines).
- An entire general fraction enclosed in general fraction indicators (⠠⠠) and (⠠⠠).
Note that Grade 1 indicators will be required if the sequence is not already in Grade 1 mode.
- An entire radical expression enclosed in radical indicators (such as a square root).
- An arrow.
- An arbitrary shape.
- Any expression enclosed in matching pairs of round parentheses, square brackets, or curly braces.

Notes:

- If none of the above conditions apply, then the “item” is the only the next symbol and may require braille grouping indicators, opening (⠠⠠) and closing (⠠⠠), to

ensure the whole of the superscript or subscript has been captured. Note that Grade 1 indicators will be required for these signs if the sequence is not already in Grade 1 mode.

- Any expression when enclosed in the braille grouping indicators (described below) will subsequently make it clear to the reader exactly what symbols are included as part of the level change.
- The superscript (⠠⠠) and subscript (⠡⠠) signs each have a Grade 2 (contracted) meaning so will always require Grade 1 indicators if the sequence is not already shown in Grade 1 mode.
- A negative superscript or subscript must be enclosed in braille grouping signs because a minus sign can be an item in its own right.
- The numeric indicator also sets Grade 1 mode for the remainder of the symbols-sequence.

Braille Grouping Symbols

Refer to the definition of an item above to determine if the braille grouping symbols will be required.

⠠ Opening braille grouping sign

⠡ Closing braille grouping sign

Bars

Common modifiers such as bars are treated separately from something that is written directly above or directly below a term.

⠠ bar over previous item

⠡ bar under previous item

Note:

- The bar directly over a previous item (⠠) and the bar directly under a previous item (⠡) have a Grade 2 (contracted) meaning, so unless you are already in Grade 1 mode, a Grade 1 symbol indicator will be required before the bar over and bar under indicators.

- Use the Grade 1 **word** indicator when there are two or more symbols in the sequence requiring Grade 1 mode except in a context whereby any literary elements will be affected and as a consequence will also be uncontracted.
- Use Grade 1 **passage** indicator (with Grade 1 terminator) when a comparison indicator or a space is used in the sequence and Grade 1 indicators are required on both sides of the equation. Remember however, the impact of Grade 1 mode on any literary elements.

Hint:

- The numeric indicator also sets Grade 1 mode for the remainder of the symbols-sequence.

Exercise 3

1. $z = \frac{x - \bar{x}}{s}$

2. $\bar{x} = \frac{\text{sum of scores}}{\text{number of scores}}$

3. $y^2 = -i\bar{y}$

4. $\bar{x} = \frac{25 + 35 + 45}{5}$

5. \underline{PQ}

6. $\underline{m + n}$

7. $4.1\dot{5}2\dot{5}$

8. $\dot{x} = \frac{t}{t^2 + 3} \text{ms}^{-1}$

9. $\ddot{x} = g - kv$

10. $\ddot{x} = 25e^{5t} \text{ms}^{-2}$

Review Exercise 3

1. $\log_x\left(\frac{1}{3}\right) = -1$

2. $\therefore \theta = 37^\circ 23'$

3. $\frac{n^2 - n - 6}{n^2 + 5n + 6}$

4. $\sqrt{\frac{-3c}{b-a}}$

5. $\log_a x = \frac{\log_b x}{\log_b a}$

6. $\log_7 \sqrt[4]{7}$

7. $\therefore \Delta RST \equiv \Delta XYZ$

8. $y = 2\sqrt{x} = 2x^{\frac{1}{2}}$

9. $d = \sqrt{(x_2 - x_1)^2}$

10. $\frac{\log_e x}{e^{2x+x}}$

Lesson 4 - Modifiers, Arrows and Limits

Modifiers

Modifiers directly above or below

If something is written directly above or directly below a term rather than to the right or left, use the directly above indicator or directly below indicator instead of the superscript or subscript.



Expression directly above



Expression directly below

Comparison signs are usually unspaced when they appear in an expression placed directly above or directly below the character.

However, a common modifier such as an arrow is treated similar to bars and a dot (described in Lesson 3) and has its **own** sign instead of using the indicators for “directly above” or “directly below”. This is introduced below in this lesson.

Definition of an Item

An “item” is defined as any single symbol(s) that follows immediately after the level change indicator. It is therefore important to make clear to the braille reader exactly what symbol(s) must be included as a consequence of the level change indicator.

Below is a defined list of specific conditions that are considered to be the “next item”:

- An entire number expressed in braille, i.e. the initiating numeric indicator and all succeeding symbols within the numeric mode (including any interior decimal points, commas, separator spaces, or simple numeric fraction lines).
- An entire general fraction enclosed in general fraction indicators (⠠) and (⠡). Note that Grade 1 indicators will be required if the sequence is not already in Grade 1 mode.
- An entire radical expression enclosed in radical indicators (such as a square root).
- An arrow.
- An arbitrary shape.
- Any expression enclosed in matching pairs of round parentheses, square brackets, or curly braces.

Arrows

For a complete list of Arrows, refer to the *Unified English Braille Guidelines for Technical Material* (International Council on English Braille, 2014).

The arrow indicator has a Grade 2 (contracted) meaning so unless you are already in Grade 1 mode, a Grade 1 symbol indicator will be required before the arrow indicator.

$\cdot\cdot$ arrow indicator

$\cdot\cdot\cdot$ → simple right pointing arrow (east)

$\cdot\cdot\cdot$ ← simple left pointing arrow (west)

Simple Arrows

- A simple arrow has a standard barbed tip at one end (like a print letter “v” on its side) pointing away from the shaft.
- The shaft is straight, and its length and thickness are not significant in braille.
- Simple arrows are represented by an arrow indicator as shown in the list above.
- Arrows are considered signs of comparison and should usually be spaced, with the exception of when they are written immediately below an item, as in Limit Functions described later in this lesson.

Examples:

$n \rightarrow 0$ (n tends to zero shown using Grade 1 symbol mode)

$\cdot\cdot\cdot$ $\cdot\cdot\cdot\cdot$ $\cdot\cdot\cdot$

OR

$n \rightarrow 0$ (n tends to zero shown using Grade 1 passage mode)

$\cdot\cdot\cdot\cdot\cdot\cdot$ $\cdot\cdot\cdot$ $\cdot\cdot\cdot\cdot\cdot\cdot$

$x \rightarrow \infty$ (x tends to infinity shown using Grade 1 symbol mode)

$\cdot\cdot\cdot$ $\cdot\cdot\cdot\cdot$ $\cdot\cdot\cdot$

OR

on both sides of the equation. Remember however, the impact of Grade 1 mode on any literary elements.

Hints:

- The numeric indicator also sets Grade 1 mode for the remainder of the symbols-sequence.
- Insert a space if a function name is **preceded** directly by a lower or upper-case Latin letter, with no intervening braille indicators or brackets.

Exercise 4

1. Input \rightarrow process \rightarrow output

2. $x \rightarrow y$

3. \overrightarrow{AB}

4. $\overrightarrow{AB} + \overrightarrow{CD}$

5. $\lim_{x \rightarrow 4} x^2 + 5$

6. $\lim_{x \rightarrow \infty} \frac{x}{x^2}$

7. $\lim_{g \rightarrow 1} \frac{g^2 - 1}{g - 1}$

8. $\lim_{t \rightarrow 4} \frac{t^2 - 16}{t - 4}$

9. $\lim_{x \rightarrow \infty} \frac{3x^2}{x^2 - 2x}$

10. $\lim_{h \rightarrow 0} \frac{2h^2x + hx^2}{h}$

Review Exercise 4

1. $a \rightarrow y = \ln(x + 1)$

2. $\frac{\sqrt{2x+1}}{(x-3)^3}$

3. $\frac{2x}{(x+5)^{\frac{1}{2}}}$

4. $OC = \sqrt{r^2 - \frac{x}{2}}$

5. $c^2 = a^2 + b^2 - 2ab \cos C$

6. $\frac{48p^2q^4}{6pq^3}$

7. $A = \pi(R^2 - r^2)$

8. $n = \frac{42 \sin 117^\circ}{\sin 35^\circ}$

9. $\cos C = \frac{a^2 + b^2 - c^2}{2ab}$

10. $\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n$

Example:

$$\sum_{x=1}^n x_i^2$$



Probability

Refer to the “*Unified English Braille Guidelines for Technical Material*” (International Council on English Braille, 2014) for further representations of Set Theory, Group Theory and Logic.

The following signs may be used in the topic of Probability at the level of Mathematics being presented in this training program.

⠠⠨ ⠠⠨ ∪ union (upright U shape): unspaced


⠠⠨ ⠠⠨ ∩ intersection (inverted ∩ shape): unspaced

⠠⠨ ⠠⠨ ∈ is an element of: spaced


⠠⠨ ⠠⠨ ⊂ is a subset of: spaced

Examples:

$A \cap B \subset A \cup B$ (A intersection B is a subset of A union B)



$3 \in A \cap B$ (3 is an element of A intersection B)



Probability is the study of how likely it is that something will happen.

Probability is when there is some doubt about the outcome, but the degree of doubt is different. Probability is about the doubt: it is the study of events that may or may not happen, rather than of events that will happen or that have already happened.

The probability of an outcome is defined as the ratio of the number of favourable outcomes to the number of possible outcomes, assuming that the outcomes are equally likely. Thus, the probability of P(A) of a particular result A is:

$$P(A) = \frac{\text{number of favourable outcomes}}{\text{number of possible outcomes}}$$

The probability of an event occurring is the number of ways the event can occur divided by the number of possible outcomes.

OR

$$P(\text{Event}) = \frac{\text{number of ways Event can occur}}{\text{number of possible outcomes}}$$

The probability of an event occurring is the number of ways the event can occur divided by the number of possible outcomes.

Note:

The example shown above has been brailled using Grade 1 symbol indicators where required rather than using Grade 1 word or Grade 1 passage mode due the number of literary elements in the sequence.

Venn Diagrams

Venn diagrams are a visual (or graphical) way to represent sets of data. You can name the sets with letters and talk about various properties of the sets.

Venn diagrams are an efficient tool to be used when analysing data to find the probability of events.

While it is not critical to understand the mathematics of Probability and Venn diagrams when transcribing braille, it may be helpful for the brief explanation below that was extracted from a student’s textbook:

Example:

A description of a Venn diagram is as follows:

There is an outer group in a rectangle containing all the members of a set S, an inner group positioned in a circle on the left within the rectangle containing all the members of set D, an inner group positioned in a circle on the right within the rectangle containing all the members of set F.

The area where the inner groups (circles) overlap contains the members that are in both sets D and F. The region outside the groups of D and F contain all the members of set S that are in neither set D nor set F.

The set made by combining sets D and F is called **union** of the two sets, written as $D \cup F$ or ⠠⠠⠠⠠⠠⠠⠠⠠ in braille. It consists of all the elements that are in D or in F or in both D and F.

The set made up of the elements common to sets D and F is called the **intersection** of the two sets, written as $D \cap F$ or ⠠⠠⠠⠠⠠⠠⠠⠠ in braille. The intersection contains numbers that belong to both sets.

Complementary Events

The **complement of event A** is the event “A does not occur” and can be denoted by \bar{A} .

If an experiment has n possible outcomes, m of which are associated with event A , then $(n - m)$ outcomes are associated with event \bar{A} .

Examples:

$$P(A) = \frac{m}{n}$$

⠠⠠⠠⠠⠠⠠⠠⠠ ⠠⠠⠠⠠⠠⠠⠠⠠ ⠠⠠⠠⠠⠠⠠⠠⠠ ⠠⠠⠠⠠⠠⠠⠠⠠

OR

⠠⠠⠠⠠⠠⠠⠠⠠ ⠠⠠⠠⠠⠠⠠⠠⠠ ⠠⠠⠠⠠⠠⠠⠠⠠ ⠠⠠⠠⠠⠠⠠⠠⠠

AND

$$P(\bar{A}) = \frac{n-m}{n}$$

⠠⠠⠠⠠⠠⠠⠠⠠ ⠠⠠⠠⠠⠠⠠⠠⠠ ⠠⠠⠠⠠⠠⠠⠠⠠ ⠠⠠⠠⠠⠠⠠⠠⠠ ⠠⠠⠠⠠⠠⠠⠠⠠

⠠⠠⠠⠠⠠⠠⠠⠠ ⠠⠠⠠⠠⠠⠠⠠⠠ ⠠⠠⠠⠠⠠⠠⠠⠠ ⠠⠠⠠⠠⠠⠠⠠⠠

⠠⠠⠠⠠⠠⠠⠠⠠ ⠠⠠⠠⠠⠠⠠⠠⠠ ⠠⠠⠠⠠⠠⠠⠠⠠ ⠠⠠⠠⠠⠠⠠⠠⠠


Independent Events

Two events are **independent** if the outcome of each event is not affected by the outcome of the other event. This means that the probability of both events together is the same as the probability of both events separately, and so their probabilities can be multiplied together as if they are separate successful outcomes. In other words:

Example:

If events A and B are independent, then:

$$P(A \text{ and } B) = P(A) \times P(B)$$




Dependent Events

Two events are **dependent** if the outcome of one event is affected by the outcome of the other event. This means that the probability of both events together is **not** the same as the probability of both events separately. Their probabilities cannot be simply multiplied together as if they are separate successive outcomes (as for independent events). In other words:

Example:

If events A and B are dependent, then:

$$P(A \text{ and } B) \neq P(A) \times P(B)$$



Remember:

As referred to in the General Note in Lesson 1, there is often a choice of Grade 1 indicator, with any of the options of Grade 1 mode (symbol, word, or passage) being equally correct.

Decisions about option selection are generally associated with user and transcriber preferences, or local production formatting guidelines, and include consideration for simplicity or functionality.

Therefore, for the purpose of the exercises in UEB Mathematics Online, use the following criteria for implementing Grade 1 mode:

- Use the Grade 1 **symbol** indicator when there is only one symbol in the sequence requiring a Grade 1 indicator.
- Use the Grade 1 **word** indicator when there are two or more symbols in the sequence requiring Grade 1 mode except in a context whereby any literary elements will be affected and as a consequence will also be uncontracted.
- Use Grade 1 **passage** indicator (with Grade 1 terminator) when a comparison indicator or a space is used in the sequence and Grade 1 indicators are required on both sides of the equation. Remember however, the impact of Grade 1 mode on any literary elements.

Hints:

- The numeric indicator also sets Grade 1 mode for the remainder of the symbols-sequence.
- If the braille representation for a print sequence does not fit on the line, then a dot 5 (⠆) continuation indicator placed at a logical place immediately following the last character may be required to show the braille is continuing to the next line. Usually the preferred place to break is before a comparison sign, an operation sign, or a fraction line indicator. For the purposes of this training program take the remainder of the sequence to the margin as it is recognised there may be various local formatting guidelines.
- Signs of operation and comparison are unspaced when they appear in an expression which is not on the base line such as directly above or directly below the character.

Exercise 5

1. $A = \frac{\theta}{360} \times \pi r^2$
2. $Z = \frac{x - \mu}{\sigma}$
3. $(\cos \frac{\pi}{4} - \tan \frac{\pi}{4})^2$
4. $\theta = 2n\pi \pm \cos^{-1} a$
5. $\tan(\theta + \phi) = \frac{\tan\theta + \phi}{1 - \tan\theta \tan\phi}$
6. $n(V \cup C) = n(V) + n(C)$
7. $X \cap Y = \{1, 11, 12\}$
8. $n(D \cup F) = n(C) + n(F) - n(D \cap F)$
9. $P(\bar{A} \text{ or } B) = 1$
10. $n(E \cap F) = 0$

Review Exercise 5

1. $\frac{pq}{5} \times \frac{p}{q}$
2. Roots of $P(z)$ are $\alpha, \bar{\alpha}, \beta, \bar{\beta}$
3. $\sin\theta = \frac{o}{h}$
4. $\bar{x} = \frac{\Sigma fx}{\Sigma f}$
5. $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$
6. $\lim_{\delta \rightarrow 0} \frac{\delta y}{\delta x}$
7. $\frac{y+1}{y-1} > 2$

8. $\sum_{x=1}^n (2^r x)$

9. $\sin\left(\pi - \frac{\pi}{6}\right)$

10. $\frac{1}{2} [\cos (A) + \cos (B)]$

- The numeric indicator also sets Grade 1 mode for the remainder of the symbols-sequence.
- Insert a space if a function name is **preceded** directly by a lower or upper-case Latin letter, with no intervening braille indicators or brackets.
- Insert a space if a function name is **followed** directly by a lower-case Latin letter, with no intervening braille indicators or brackets.
- If the braille representation for a print sequence does not fit on the line, then a dot 5 (⠨) continuation indicator placed at a logical place immediately following the last character may be required to show the braille is continuing to the next line. Usually the preferred place to break is before a comparison sign, an operation sign, or a fraction line indicator. For the purposes of this training program take the remainder of the sequence to the margin as it is recognised there may be various local formatting guidelines.

Exercise 6

1. $\frac{dy}{dx} > 0$

2. $\frac{dy}{dx} f'(x)$

3. $\frac{dy}{dx} f'(x) e^{f(x)}$

4. $\frac{d}{dx} (x^2 - 2x)$

5. $\frac{dy}{dx} = \frac{2}{\sqrt{1-x^2}}$

6. $\frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$

7. $\frac{d}{dx} (kx^n) = knx^{n-1}$

8. $\frac{dy}{dx} = 4x^3 - 9x^2 + 7$

9. $\frac{dy}{dx} = f'(x) \cos f(x)$

10. $\frac{d}{dx} [f(x)]^2 =$

Review Exercise 6

1. $\log_b x = \frac{\log_a x}{\log_a b}$

2. $\frac{\delta y}{\delta x} = \frac{\delta y}{\delta u} \times \frac{\delta u}{\delta x}$

3. $\frac{dy}{dx} = nx^{nx-1}$

4. $\frac{ax_1 + by_1 + c}{\sqrt{a^2 + b^2}}$

5. $\frac{dy}{dx} = \frac{f'(x)}{f(x)}$

6. $\frac{dy}{dx} = 5x^4 - 9$

7. $\left(\frac{ab^5c}{abc^4}\right)^0$

8. $\lim_{\theta \rightarrow 0} \frac{\tan \frac{\theta}{3}}{\theta}$

9. $\lim_{x \rightarrow \infty} \frac{3\sqrt{x}}{\sqrt{x} - 1}$

10. $\lim_{x \rightarrow -1} \frac{x^2 + 4x + 1}{x^2 - 1}$

Lesson 7 - Calculus: Integration

As Calculus is a defined topic in senior secondary level mathematics it has been introduced specifically into the lessons for this training program. The same rules and principles of braille transcription should be observed that have been introduced previously.

There are two main branches of Calculus: Differentiation and Integration. The topic of Differentiation was introduced in Lesson 6.

Integration as presented in this Lesson is the inverse of differentiation and uses information about rates of change to go back and examine the original variables. Integration can also be used to find areas of curved objects.

This training program will only be introducing the standard integral sign (\int) at the level of Mathematics being presented in this training program and is shown in braille as (\int).

Notes:

- The spacing of the integral sign in print (\int) may be unclear or inconsistent. In braille it is best to have the integral sign (\int) unspaced from the function and treat its limits as subscripts and superscripts. The dx at the end means “integrate with respect to x ” and can also be written unspaced.
- While both of the examples below are correct, the first example in each using Grade 1 word mode is the preferred option due to requiring more than one (1) Grade 1 symbol indicator in the sequence.

Examples:

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx$$

$\int \frac{1}{\sqrt{a^2 - x^2}} dx$ preferred

OR

$\int \frac{1}{\sqrt{a^2 - x^2}} dx$

$$\int_2^3 (2x - 1) dx$$

$\int_2^3 (2x - 1) dx$ preferred

OR

$\int_2^3 (2x - 1) dx$

Remember:

As referred to in the General Note in Lesson 1, there is often a choice of Grade 1 indicator, with any of the options of Grade 1 mode (symbol, word, or passage) being equally correct.

Decisions about option selection are generally associated with user and transcriber preferences, or local production formatting guidelines, and include consideration for simplicity or functionality.

Therefore, for the purpose of the exercises in UEB Mathematics Online, use the following criteria for implementing Grade 1 mode:

- Use the Grade 1 **symbol** indicator when there is only one symbol in the sequence requiring a Grade 1 indicator.
- Use the Grade 1 **word** indicator when there are two or more symbols in the sequence requiring Grade 1 mode except in a context whereby any literary elements will be affected and as a consequence will also be uncontracted.
- Use Grade 1 **passage** indicator (with Grade 1 terminator) when a comparison indicator or a space is used in the sequence and Grade 1 indicators are required on both sides of the equation. Remember however, the impact of Grade 1 mode on any literary elements.

Hints:

- The numeric indicator also sets Grade 1 mode for the remainder of the symbols-sequence.
- Insert a space if a function name is **preceded** directly by a lower or upper-case Latin letter, with no intervening braille indicators or brackets.
- Insert a space if a function name is **followed** directly by a lower-case Latin letter, with no intervening braille indicators or brackets.
- Signs of operation and comparison are unspaced when they appear in an expression which is not on the base line such as directly above or directly below the character.
- If the braille representation for a print sequence does not fit on the line, then a dot 5 (⠨) continuation indicator placed at a logical place immediately following the last character may be required to show the braille is continuing to the next line. Usually the preferred place to break is before a comparison sign, an operation sign, or a fraction line indicator. For the purposes of this training program take the remainder of the sequence to the margin, as it is recognised there may be various local formatting guidelines.

Exercise 7

1. $\int x^2 dx$

2. $\int_0^2 \frac{3}{4+x^2} dx$

3. $\int 2x^4 dx$

4. $\int_3^\pi \cos\left(\frac{x}{2} + \pi\right) dx$

5. $\int_0^5 \frac{dx}{x+3}$

6. $\int_a^b f(x) dx$

7. $\int \sin^2 x dx$

8. $\int e^{ax+b} dx$

9. $\int \tan x dx = \ln|\cos x| + c$

10. $\pi \int_0^{\frac{\pi}{4}} \sec^2 x dx$

Review Exercise 7

1. $f''(x) = 6x - 8$

2. $\alpha\beta + \alpha\gamma + \beta\gamma = \frac{c}{a}$

3. $\frac{d^2x}{dt^2}$

4. $P(E') = 1 - p(E)$

5. $\frac{dy}{dx} = F'(u) \frac{du}{dx}$

6. $\int \sec^2(ax + b) dx$

7. $\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$

8. $\int \cos(ax + b) dx$

9. $\int_1^2 \frac{x^3 - x^2 + 5x + 3}{x} dx$

10. $\int (ax + b)^n$

Exercise 8

1. $|y - 4|$

2. $\frac{|x|}{x^2} = 1$

3. $|x| \geq 0$

4. $\sqrt{a^2} = |a|$

5. $\frac{|3x - 2|}{3x - 2} = 1$

6. $|3y - 1| + |2y + 3| > 5$

7. $y^2 = 2\sin|x|$

8. $|z - 2| + |z + 1| = 7$

9. $|3x + 2| = 2x - 3$

10. $|\overrightarrow{O_1P}|$

Review Exercise 8

1. $A = \frac{h}{2}(d_f + d_1)$

2. $(x - h)^2 = \pm 4a(y - k)$

3. $y = e^{f(x)}$

4. $\frac{p^{\frac{1}{2}} \times p^{\frac{1}{2}}}{p^2}$

5. $S_n = \frac{a(r^n - 1)}{r - 1}$

6. $x^{\frac{1}{n}} = \sqrt[n]{x}$

7. $\frac{1}{\sqrt[4]{(7-x)^2}}$

8. $\left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$

9. $f'(x)n[f(x)]^{n-1}$

10. $\lim_{h \rightarrow 0} \frac{2xh - h^2 - 5h}{h}$

Lesson 9 - Review Test

Congratulations on reaching this Review Test. The content of each lesson has built upon preceding lessons, with the overall structure of the training manual designed to reinforce several foundational principles, including the following:

1. The numeric indicator sets numeric mode **and** Grade 1 mode for the remainder of the symbols-sequence unless terminated for any reason.
2. A braille symbol may have both an uncontracted (Grade 1) meaning and a contracted (Grade 2) meaning. A Grade 1 indicator is used to set Grade 1 mode when the Grade 1 meaning of a symbol could be misread as a contraction meaning or a numeric meaning.
3. When is a fraction considered a “simple fraction”? Note, if the sequence does not satisfy the definition of a simple fraction, then it must be treated as a general fraction.
4. The rules associated with “the next item” should be considered and are particularly relevant in the use of level change indicators.

Remember:

As referred to in the General Note in Lesson 1 in this UEB Training Manual: Extension Mathematics and throughout the lessons, the choice of options for Grade 1 mode (symbol, word, or passage) may be equally correct in mathematical contexts.

Decisions about option selection are generally associated with user and transcriber preferences, or local production formatting guidelines, and include consideration for simplicity or functionality.

Therefore, for the purpose of the exercises in UEB mathematics Online, use the following criteria for implementing Grade 1 mode:

- Use the Grade 1 **symbol** indicator when there is only one symbol in the sequence requiring a Grade 1 indicator.
- Use the Grade 1 **word** indicator when there are two or more symbols in the sequence requiring Grade 1 mode except in a context whereby any literary elements will be affected and as a consequence will also be uncontracted.
- Use Grade 1 **passage** indicator (with Grade 1 terminator) when a comparison indicator or a space is used in the sequence and Grade 1 indicators are required on both sides of the equation. Remember however, the impact of Grade 1 mode on any literary elements.

Hints:

- The numeric indicator also sets Grade 1 mode for the remainder of the symbols-sequence.

- Signs of operation and comparison are unspaced when they appear in an expression which is not on the base line such as directly above or directly below the character.
- If the braille representation for a print sequence does not fit on the line, then a dot 5 (⠄) continuation indicator placed at a logical place immediately following the last character may be required to show the braille is continuing to the next line. Usually the preferred place to break is before a comparison sign, an operation sign, or a fraction line indicator. For the purposes of this training program take the remainder of the sequence to the margin, as it is recognised there may be various local formatting guidelines.

Review Test

1. $8.754\dot{7}5\dot{9}$

2. $T_n = a + (n - 1)$

3. $\bar{y} = -10$

4. \overline{AB}

5. $\ln |f(x)| + c$

6. $\bar{x} = \frac{\Sigma x}{n}$

7. $|5x - 3| \geq 2$

8. $\frac{(2x^7)^3 y^2}{x^{10} y}$

9. $[2(x + y)] \div 4 < 10$

10. $|-8| = |8| = 8$

11. $\cos C = \frac{a^2 + b^2 - c^2}{2a}$

12. $A = \int_{-1}^2 (2 + x) dx$

13. $\int_b^c f(x) dx$

14. $\sin(\pi - \frac{\pi}{8})$

15. $\frac{n!}{r!(n-r)!}$

16. $\theta = n\pi + (-1)^n \sin^{-1} a$

17. $\lim_{x \rightarrow 0} f(x)$

18. $f_1(x) + f_2(x)$

19. $S_n = \frac{n}{2}[2a + (a - l)d]$

20. $f''(x) = k(b^2 - x^2)$

Lesson 10 - Extension Test

Congratulations on reaching this Extension Test. As noted in Lesson 9, the content of each lesson has built upon preceding lessons, with the overall structure of the training manual designed to reinforce several foundational principles, including the following:

1. The numeric indicator sets numeric mode **and** Grade 1 mode for the remainder of the symbols-sequence unless terminated for any reason.
2. A braille symbol may have both an uncontracted (Grade 1) meaning and a contracted (Grade 2) meaning. A Grade 1 indicator is used to set Grade 1 mode when the Grade 1 meaning of a symbol could be misread as a contraction meaning or a numeric meaning.
3. When is a fraction considered a “simple fraction”? Note: If the sequence does not satisfy the definition of a simple fraction, then it must be treated as a general fraction.
4. The rules associated with “the next item” should be considered and are particularly relevant in the use of level change indicators.

Remember:

As referred to in the General Note in Lesson 1 in this UEB Training Manual: Extension Mathematics and throughout the previous lessons, the choice of options for Grade 1 mode (symbol, word, or passage) may be equally correct in mathematical contexts.

Decisions about option selection are generally associated with user and transcriber preferences, or local production formatting guidelines, and include consideration for simplicity or functionality.

Therefore, for the purpose of the Extension Test in this UEB Training Manual, use the following criteria for implementing Grade 1 mode:

- Use the Grade 1 **symbol** indicator when there is only one symbol in the sequence requiring a Grade 1 indicator.
- Use the Grade 1 **word** indicator when there are two or more symbols in the sequence requiring Grade 1 mode except in a context whereby any literary elements will be affected and as a consequence will also be uncontracted.
- Use Grade 1 **passage** indicator (with Grade 1 terminator) when a comparison indicator or a space is used in the sequence and Grade 1 indicators are required on both sides of the equation. Remember however, the impact of Grade 1 mode on any literary elements.

Hints:

- The numeric indicator also sets Grade 1 mode for the remainder of the symbols-sequence.

- Insert a space if a function name is **preceded** directly by a lower or upper-case Latin letter, with no intervening braille indicators or brackets.
- Insert a space if a function name is **followed** directly by a lower-case Latin letter, with no intervening braille indicators or brackets.
- Signs of operation and comparison are unspaced when they appear in an expression which is not on the base line such as directly above or directly below the character.
- If the braille representation for a print sequence does not fit on the line, then a dot 5 (⠨) continuation indicator placed at a logical place immediately following the last character may be required to show the braille is continuing to the next line. Usually the preferred place to break is before a comparison sign, an operation sign, or a fraction line indicator. For the purposes of this training program take the remainder of the sequence to the margin as it is recognised there may be various local formatting guidelines.

Extension Test

1. $x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}$

2. $\int x^2 dx$

3. $\int \frac{\cos x}{\sin x} dx$

4. $\int_2^3 \frac{dx}{x}$

5. $d = \sqrt{(x_2 - x_1)}$

6. $\operatorname{cosec} A = \frac{1}{\sin A}, \sin A \neq 0$

7. $(\operatorname{Re}(\alpha))^2 + (\operatorname{Re}(\beta))^2 = 1$

8. $\int \frac{6x^2}{x^3}$

9. $d = \frac{|ax_1|}{\sqrt{a^2}}$

10. $\frac{d}{dx} \left(\frac{x^2}{x-1} \right)$

11. $\bar{x} = \frac{\sum x}{n}$

12. $\sum_2^4 |3p - 2|$

13. $\ddot{x} = -n^2(x - c)$

14. $y = \tan^{-1} f(x)$

15. $A_n = P \left(1 + \frac{r}{100} \right)^n$

16. $\left(\frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n} \right)$

17. $\max_{0 \leq x \leq 1} x e^{-x^2}$

18. $\sum_{i=1}^n x_i p_i$

19. $\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n$

20. $\int \frac{f'(x)}{f(x)} dx$

References

- Howse, J. (2018a). Unified English Braille Maths K – 6. Workshop presented at RIDBC Renwick Centre, North Rocks, NSW. Unpublished document.
- Howse, J. (2018b). Unified English Braille Maths 7 – 12 Workshop presented at RIDBC Renwick Centre, North Rocks, NSW. Unpublished document.
- Howse, J. (2019a). Unified English Braille training manual: Introductory mathematics (Revision 5). <https://uebonline.org/resources/ueb-mathematics-resources/>
- Howse, J. (2019b). Unified English Braille training manual: Advanced Mathematics (Revision 3). <https://uebonline.org/resources/ueb-mathematics-resources/>
- Howse, J. Riessen, K. and Holloway, L. (Eds.). (2016). Unified English Braille: Australian Training Manual. <http://brailleaustralia.org/unified-english-braille/unified-english-braille-australian-training-manual-2013/>
- International Council on English Braille. (2014). Unified English Braille Guidelines for Technical Material. http://www.iceb.org/guidelines_for_technical_material_2014.pdf
- Simpson, C. (Ed.). (2013).
- The Rules of Unified English Braille (2013) (2nd ed.). International Council on English Braille (ICEB). [http://iceb.org/Rules of Unified English Braille 2013.pdf](http://iceb.org/Rules_of_Unified_Engish_Braille_2013.pdf).